

Some questions about the dimension of a group action.

Collin Bleak

Department of Mathematics, Cornell University, Ithaca, NY 14853-4201 USA

Abstract

We discuss three families of groups, ZW_n , $PL(I^n)$, and $PL(S^n)$ (the last two being families of groups of piecewise-linear homeomorphisms of standard n -dimensional spaces). We show that for positive $n \in \mathbf{N}$, ZW_n embeds in $PL(I^n)$ which embeds in $PL(S^n)$. In another direction, ZW_2 fails to embed in $PL(I^1)$ by a result in [2]. Here, we extend that result to show that ZW_2 also fails to embed in $PL(S^1)$. The nature of the proofs of these non-embedding results leads us to ask if there are corresponding non-embedding results in higher dimensions.

Contents

1	Introduction	2
1.1	Families of groups of piecewise-linear self-homeomorphisms	2
1.2	Standard restricted wreath products, and another family of groups	2
1.3	Notation and theory associated with group actions and homeomorphisms of the circle	3
1.4	Statements of Results	4
1.5	Two Questions	4
2	ZW_2 fails to embed in $PL(S^1)$	5

1 Introduction

We use a non-embedding result in [2] to show that the group $\mathbf{Z} \wr (\mathbf{Z} \times \mathbf{Z})$ does not embed in $PL(S^1)$, the group of piecewise-linear homeomorphisms of S^1 (under the operation of composition). This, together with a family of easy embedding results for a family ZW_n of related groups leads us to ask some natural questions about the parallel results in higher dimensions.

The author would like to thank Ken Brown for asking whether the non-embedding of $\mathbf{Z} \wr (\mathbf{Z} \times \mathbf{Z})$ into $PL_o(I)$ (and therefore into R. J. Thompson's group F) could be extended to a non-embedding of $\mathbf{Z} \wr (\mathbf{Z} \times \mathbf{Z})$ into Thompson's group T , a subgroup of $PL(S^1)$. The author is also indebted to Martin Kassabov and Francesco Matucci for simplifying the arguments below in accord with our on-going joint work (see [3]).

1.1 Families of groups of piecewise-linear self-homeomorphisms

For each positive integer n , let I_p^n and S_p^n represent polyhedra in $\mathbf{R}^n$ and $\mathbf{R}^{n+1}$ which are homeomorphic to I^n and S^n respectively, where we further require that $\mathbf{R}^{n+1}$ admits a reflective plane of symmetry for the polyhedron S_p^n , so that S_p^n is realized as the union of two piecewise-linear homeomorphic images of I_p^n , identified together on their common boundary, where one image is the reflection of the other across the plane of symmetry, excepting their boundaries, which lie in the plane of symmetry itself. We will refer to $\text{Homeo}(I^n)$ and $\text{Homeo}(S^n)$ as the full groups of self-homeomorphisms of the polyhedra I_p^n and S_p^n respectively, under composition, and we will refer to $PL(I^n)$ and $PL(S^n)$ as the subgroups of these full homeomorphism groups consisting of the piecewise-linear self-homeomorphisms.

1.2 Standard restricted wreath products, and another family of groups

We will be discussing a third family of groups as well. These groups are standard restricted wreath products of well known groups. Wreath products are discussed in detail in many places, but two standard references are Peter Neumann's paper [7] and the book [6] by J. D. P. Meldrum. We now give a general description of the standard restricted wreath product.

Let the symbol $C \wr T$ represent the standard restricted wreath product of the groups C and T . In particular, $C \wr T \cong B \rtimes T$, where $B \cong \bigoplus_{t \in T} C$. We use the coordinate system provided by the isomorphism of $C \wr T$ with $B \rtimes T$ to describe elements of $C \wr T$. In particular, if $((a_t)_{t \in T}, r), ((b_t)_{t \in T}, s) \in C \wr T$, then we describe the product of these two elements by the following formula.

$$((a_t)_{t \in T}, r) \cdot ((b_t)_{t \in T}, s) = ((c_t)_{t \in T}, rs) \text{ where for } t \in T, \text{ we have } c_t = a_t b_{(t,r)}.$$

In this notation, C is called the bottom group, B is called the base group, and T is called the top group. We often think of the base and top groups as being identified with the standard appropriate subgroups in $B \rtimes T$, and we will sometimes refer to B and T as subgroups of $C \wr T$.

Given any positive integer n , let ZW_n represent the group $\mathbf{Z} \wr (\mathbf{Z} \times \mathbf{Z} \times \dots \times \mathbf{Z})$, where there are n appearances of the factor $\mathbf{Z}$ in the direct product.

1.3 Notation and theory associated with group actions and homeomorphisms of the circle

All group actions in this paper will be right actions, that is, if x is a point in a space S acted on by a group G and $a \in G$, then we will denote by xa the image of x under the action of a . Likewise, if $X \subset S$, we will denote by Xa the set $\{xa \mid x \in X\}$. Consistent that notation, if b is also in G , we will use the following notational conventions, $a^b = b^{-1}ab$ and $[a, b] = a^{-1}b^{-1}ab$. Standard to this variety of notation is the equation $\text{Support}(a^b) = (\text{Support}(a))b$, where the support of an element of G will be the set of points in S moved by the element.

There are three main results that we will use in our arguments, that are relevant to the classic body of theory associated with groups of homeomorphisms of the circle.

The first result is the classical Ping-Pong Lemma, the proof of which can be found in many places. The earliest reference I know of is [5], a paper in German by Felix Klein. The statement I give follows that in [1].

Lemma 1.1 (Ping-Pong Lemma). *Let Γ be a group of permutations on a set X , let g_1, g_2 be elements of Γ of order at least three. If X_1 and X_2 are disjoint subsets of X and for all $n \neq 0, i \neq j, X_j g_i^n \subset X_i$, then g_1, g_2 freely generate the free group F_2 .*

While this lemma is clearly not restricted to groups acting on the circle, it plays a heavy role in the theory of such groups, particularly when the action of such a group is highly transitive.

We will also make heavy use of Poincaré's concept of the rotation number of a self-homeomorphism of S^1 .

Given a homeomorphism $f : S^1 \rightarrow S^1$, let $F : \mathbf{R} \rightarrow \mathbf{R}$ represent a lift of this homeomorphism via the standard covering projection $\exp : \mathbf{R} \rightarrow S^1$, so that $0 \leq F(0) < 1$, and where we think of S^1 as a subset of the complex plane, and use $\exp(t) = e^{2\pi it}$. We then define

$$\text{Rot}(f) = \lim_{n \rightarrow \infty} \frac{F^n(x) - x}{n}.$$

Poincaré showed that this limit is well defined and it is independent of the initial point $x \in [0, 1)$ used. Given an $f \in \text{Homeo}(S^1)$, we call $\text{Rot}(f)$ the *rotation number of f* . If H is an abelian subgroup of $\text{Homeo}(S^1)$, then the rotation number map represents a homomorphism from H to the additive group $\mathbf{R} \bmod (1)$.

The second result we will need is the following lemma proved by Poincaré circa 1900.

Lemma 1.2 (Poincaré's Lemma). *Suppose $f : S^1 \rightarrow S^1$ is a homeomorphism with rational rotation number $\frac{r}{s}$ (in lowest terms, use $r = 0, s = 1$ if the rotation number is zero), then f^s has a fixed point.*

The final result is a theorem by Denjoy, although we use an extension given by M. Herman on page 76 of [4].

Theorem 1.3 (Denjoy's Theorem). *Suppose $h : S^1 \rightarrow S^1$ is either a piecewise-linear homeomorphism or a C^1 diffeomorphism whose derivative has bounded variation. If the rotation number of h is irrational, then h is conjugate (by an element in $\text{Homeo}(S^1)$) to a rotation.*

1.4 Statements of Results

In [2], we see that ZW_2 does not embed in $PL_o(I)$ (the subgroup of $PL(I^1)$ consisting only of orientation-preserving elements), and therefore, that ZW_2 does not embed in R. Thompson's group F , answering a question of Mark Sapir. In a discussion of that work, Ken Brown asked the author whether it could be extended to get a similar result for R. Thompson's group T . We in fact show the following theorem.

Theorem 1.4. *ZW_2 does not embed in $PL(S^1)$.*

Since T is often realized as a subgroup of $PL(S^1)$, we immediately obtain the following result.

Corollary 1.5. *ZW_2 does not embed in R. Thompson's group T .*

Taking the above together with the following easy result (which we leave for the reader's entertainment), we are led to ask (in the next sub-section) some natural follow-up questions.

Theorem 1.6. *Given any natural number $n > 0$, ZW_n embeds in $PL(I^n)$, and also in $PL(S^n)$.*

1.5 Two Questions

The chief non-embedding result of [2] shows that ZW_2 fails to embed in $PL_o(I)$, which immediately implies that ZW_2 fails to embed in $PL(I^1)$. The chief non-embedding result here shows that ZW_2 fails to embed in $PL(S^1)$. The nature of the proof of the non-embedding of ZW_2 in $PL(I^1)$ leads the author to suspect that these non-embedding results will generalize to higher dimensions, but he has no idea as to how to approach any proof of that suspicion. Therefore, due to a lack of large quantities of creditable evidence, we do not put forth any conjectures, but instead we ask some questions.

Question 1. *Does there exist a natural number $n > 1$ so that ZW_n embeds in $PL(I^{n-1})$?*

Note that given any piecewise-linear homeomorphism of the unit n -cube, it is easy to build an induced piecewise linear homeomorphism on the n -sphere by using the n -cube homeomorphism on both halves of the sphere, which we recall is constructed as a doubled n -cube. In particular, we have the following remark.

Remark 1.7. *For all $n \in \mathbf{N}$, the group $PL(I^n)$ embeds in $PL(S^n)$.*

The remark still leaves open the possibility that there is an $n \in \mathbf{N}$ so that ZW_n embeds in $PL(S^{n-1})$, even if it does not embed in $PL(I^{n-1})$. Therefore, the following question is of independent interest.

Question 2. *Does there exist $n \in \mathbf{N}$ so that ZW_n embeds in $PL(S^{n-1})$?*

Therefore, we have the following diagram, characterizing the situation, where the symbols $PL(A^n)$ can be taken to represent either $PL(I^n)$ or $PL(S^n)$.

$$\begin{array}{ccc}
& \vdots & \\
ZW_4 & \xhookrightarrow{\quad} & PL(A^4) \\
& \searrow ? & \\
ZW_3 & \xhookrightarrow{\quad} & PL(A^3) \\
& \searrow ? & \\
ZW_2 & \xhookrightarrow{\quad} & PL(A^2) \\
& \searrow \exists & \\
ZW_1 & \xhookrightarrow{\quad} & PL(A^1)
\end{array}$$

2 ZW_2 fails to embed in $PL(S^1)$

The proof which follows can be replaced by another proof, which relies on the classification of the solvable subgroups of $PL(S^1)$ which is forthcoming in [3]. However, the proof below is elementary, only relying on the theory which was mentioned in the introduction.

Proof of Theorem 1.4

Suppose there exists an injective homomorphism $\phi : ZW_2 \rightarrow PL(S^1)$. Recall that $ZW_2 = \mathbf{Z} \wr (\mathbf{Z} \times \mathbf{Z}) = \mathbf{Z} \wr (\mathbf{Z}^2)$. Choose $a = ((0)_{* \in \mathbf{Z}^2}, (1, 0))$ and $b = ((0)_{* \in \mathbf{Z}^2}, (0, 1))$ as generators of the top group $\mathbf{Z}^2$ in ZW_2 , and let us use $c = ((c_*)_{* \in \mathbf{Z}^2}, (0, 0))$, (where $c_* = 1$ if $* = (0, 0)$ and $c_* = 0$ otherwise) as our generator for an embedded copy of the bottom group $\mathbf{Z}$ into the base group in ZW_2 , located at index $(0, 0)$ in the base group.

Let $a\phi = \alpha$, $b\phi = \beta$, and $c\phi = \gamma$. Our argument will break into cases.

Case 1: α or β has a rational rotation number.

We will assume without meaningful loss of generality that α has a rational rotation number. If this number is not zero, then by Poincaré's Lemma, there is an integer n so that α^n has a fixed point. Further, as α has infinite order, we need not be concerned that α^n is trivial. Replace a by a^n . The new a and the old b still generate a group isomorphic to $\mathbf{Z}^2$, and $\langle a, b, c \rangle$ is still isomorphic with ZW_2 . Therefore, we may assume that the rotation number of α is zero, and that α is non-trivial, but still admits a fixed point p .

Let r be any piecewise-linear homeomorphism of S^1 which sends p to $1 \in S^1$, so that $\alpha^r = r^{-1}\alpha r$ will be a piecewise-linear homeomorphism which fixes 1. We can now replace ϕ by a new embedding of ZW_2 into $PL(S^1)$ defined by the rule $f \mapsto (\phi(f))^r$ for all $f \in ZW_2$. From now on, when we refer to ϕ , we will mean the new version of ϕ , and when we refer to α , β , or γ , we will be referring to the images of a , b , and c under this new version of ϕ .

Choose the representative lift $\tilde{\alpha} : \mathbf{R} \rightarrow \mathbf{R}$ of α so that $0\tilde{\alpha} = 0$ (and therefore $1\tilde{\alpha} = 1$, etc.). We may also assume that we chose p intelligently, so that $\tilde{\alpha}$ has a component of support $\tilde{A}$ of the form $(x, 1)$ for some $x \in (0, 1)$. Let $A = \exp(\tilde{A})$ be the image of $\tilde{A}$ under the covering projection $\exp : \mathbf{R} \rightarrow S^1$. Now, if β has an irrational rotation number, by the extended

version of Denjoy's theorem, we can find a natural number n so that $1\beta^n = y \in A$. The point y represents a fixed point of the map $\alpha^{(\beta^n)}$, where α has no fixed point. In this case, we see that α and β fail to commute. Hence β must have a rational rotation number, and in fact, some power β^k of β must fix the point 1. Replace b by b^k , ZW_2 by its isomorphic subgroup generated by a , the new b , and c , and replace β by the image of the new b under ϕ .

Now we have that α and β fix the point $1 \in S^1$, although both elements have non-trivial components of support. We also understand that if K is a component of support of α , then it is also a component of support of β , or else it is completely disjoint from the support of β , and vice-versa.

Note that $[c, a] = c^{-1}a^{-1}ca$ is a non-trivial element in the base group of ZW_2 , and it has the property that $[c, a]$, a , and b generate a subgroup of ZW_2 which is isomorphic to ZW_2 . Let θ be the image of $[c, a]$ under ϕ . The element θ is the product of two elements with fixed points, so if θ does not have fixed points, then we must have that α and $\alpha^\gamma = \phi(c^{-1}a^{-1}c)$ generate a group which contains a free group on two generators (by Klein's classical ping-pong argument (the earliest reference I know of is [5]), where we use small neighborhoods of the ends of the components of the support of the two elements to build the two required sets for the "Ping-pong"). In particular, as ZW_2 does not contain a free group on two generators, we see that θ must admit fixed points. We now replace c by $[c, a]$, so that our new γ is the θ discussed in this paragraph.

If the set of elements $\Gamma = \{\alpha, \beta, \gamma\}$ admits a global fixed point, then it is easy to see that we can embed the group $\langle \Gamma \rangle \cong ZW_2$ into $PL(I^1)$, which we know is impossible, by the chief non-embedding result of [2]. If Γ does not admit a non-empty global fixed set, then we can find an element θ in $\langle \alpha, \beta \rangle$ whose fixed set is entirely contained in the support of γ , so that again a ping-pong argument shows that ZW_2 must contain an embedded copy of the free group on two generators.

Hence, we have found contradictions in all cases, so that if ZW_2 embeds in $PL(S^1)$, the generators a and b must be realized by elements with irrational rotation numbers.

Case 2: γ has a rational rotation number.

We immediately replace c by some non-trivial power of itself so that γ has a fixed point, and we let p represent the end of some component of support D of γ .

Since α must have an irrational rotation number, the orbit of p under the action of α is dense in S^1 , and so there is an integer power k so that $p\alpha^k \in D$. But now the elements γ and γ^{α^k} cannot commute with each other, which contradicts the fact that they are both in the image of the base group of ZW_2 under ϕ , which group is abelian.

Taking this contradiction together with the results of the previous case, we see that if ZW_2 embeds in $PL(S^1)$, we must have that all of the generators α, β, γ of the image have irrational rotation number. Further, observe that no non-trivial element ω of the base group can have image with rational rotation number, or the group $\langle \alpha, \beta, \omega \rangle \cong ZW_2$ where the generator of the new bottom group would have rational rotation number.

Case 3: None of α, β , or γ has a rational rotation number.

Apply the extended version of Denjoy's theorem to find $\theta \in \text{Homeo}(S^1)$ so that α^θ is

a pure rotation. Replace α , β and γ by their images under conjugation by θ , so that the group they generate is now a subgroup of $Homeo(S^1)$ (note that the image of γ under this conjugation may not be piecewise linear).

If γ is also a pure rotation, then α and γ would commute. In particular, there is a power n of α so that the graphs of α^{-n} and γ will intersect each other on the torus (consider that the graphs of the powers of α produce a dense subset of the foliation of the torus by the graphs of the pure rotations). In particular, the functions $\alpha^{-n}\gamma^{-1}$ and $\alpha^n\gamma$ both have non-trivial fixed sets.

If the product $[\alpha^n, \gamma] = \alpha^{-n}\gamma^{-1}\alpha^n\gamma$ has no fixed points, then we see that the fixed set of $\alpha^{-n}\gamma^{-1}$ must be contained in the support of $\alpha^n\gamma$, and vice versa, so that one can use a ping-pong argument to show that ZW_2 contains a free group on two generators.

If $[\alpha^n, \gamma]$ has some fixed points, then we have found a non-trivial element in the base group of ZW_2 with rational rotation number, which is impossible by the work in our previous case.

◇

References

- [1] R. Alperin, B. Farb, and G. Noskov, *A strong schottky lemma for nonpositively curved singular spaces*, Geometriae Dedicata **92** (2002), 235–243.
- [2] Collin Bleak, *A geometric classification of some solvable groups of homeomorphisms*, preprint (2006), 24 pps.
- [3] Collin Bleak, Martin Kassabov, and Francesco Matucci, *Structure theorems for subgroups of homeomorphism groups*, in preparation (2006).
- [4] M. Herman, *Sur la conjugaison différentiable des difféomorphismes du cercle à des rotations*, Inst. Études Sci. Publ. Math. **49** (1979).
- [5] F. Klein, *Neue beitrage zur riemannischen funktionentheorie*, Math. Ann. **21** (1883), 141–218.
- [6] J. D. P. Meldrum, *Wreath products of groups and semigroups*, Pitman Monographs and Surveys in Pure and Applied Mathematics, vol. 74, Longman, Harlow, 1995. MR MR1379113 (97j:20030)
- [7] Peter M. Neumann, *On the structure of standard wreath products of groups*, Math. Z. **84** (1964), 343–373.